

AMENDMENTS TO THE CLAIMS

Claims:

1. (Currently amended) A recombinant HuEPO-L-vFc fusion protein ~~comprising~~ consisting of HuEPO, a peptide linker, and a human IgG Fc variant, wherein the human IgG Fc variant comprises a hinge, CH2, and CH3 domains of human IgG2 with Pro331Ser mutation as SEQ ID NO: 18.
2. (Original) The peptide linker in claim 1 containing about 20 or fewer amino acids is present between HuEPO and the human IgG Fc variant; and the peptide linker comprises two or more amino acids selected from the group consisting of glycine, serine, alanine, and threonine.
3. (Withdrawn) ~~The human IgG Fc variant in claim 1 comprising a hinge, CH2, and CH3 domains of human IgG2 with Pro331Ser mutation as SEQ ID NO: 18.~~
4. (Withdrawn)
5. (Withdrawn)
6. (Original) The HuEPO-L-vFc fusion protein of claim 1 which exhibits an enhanced *in vitro* biological activity of at least 2 fold relative to that of rHuEPO on a molar basis.
7. (Previously amended) A CHO cell line transfected with DNA encoding the recombinant HuEPO-L-vFc fusion protein of claim 1 in its growth medium in excess of 10 µg per million cells in a 24 hour period.
8. (Previously amended) The CHO cell line transfected with DNA encoding the recombinant HuEPO-L-vFc fusion protein of claim 7 in its growth medium in excess of 30 µg per million cells in a 24 hour period.

9. (Previously Amended) The CHO cell line transfected with DNA encoding the recombinant HuEPO-L-vFc fusion protein of claim 1, wherein the human IgG Fc variant comprises a hinge, CH2, and CH3 domains of human IgG selected from the group consisting of IgG1 as SEQ ID NO: 22, IgG2 as SEQ ID NO: 18, and IgG4 as SEQ ID NO: 20, the IgG Fc contains amino acid mutations to attenuate effector functions, a flexible peptide linker containing about 20 or fewer amino acids is present between HuEPO and human IgG Fc variant, and the HuEPO-L-vFc fusion protein exhibits an enhanced *in vitro* biological activity of at least 2 fold relative to that of rHuEPO on a molar basis.
10. (Currently Amended) A method for making a recombinant fusion protein ~~comprising~~ consisting of HuEPO, a flexible peptide linker, and a human IgG Fc variant, wherein the human IgG Fc variant comprises a hinge, CH2, and CH3 domains of human IgG2 with Pro331Ser mutation as SEQ ID NO: 18, which method comprises: (a) generating a CHO cell line transfected with DNA encoding the recombinant HuEPO-L-vFc fusion protein; (b) growing the cell line under conditions the recombinant protein is expressed in its growth medium in excess of 10 µg per million cells in a 24 hour period; and (c) purifying the expressed protein from step (b), wherein the recombinant fusion protein exhibits an enhanced *in vitro* biological activity of at least 2 fold relative to that of rHuEPO on a molar basis.
11. (Original) The method of claim 10, wherein step (b) is in excess of 30 µg per million cells in a 24 hour period.
12. (Original) The method of claim 10, wherein the flexible peptide linker containing about 20 or fewer amino acids is present between HuEPO and the human IgG Fc variant; and the peptide linker comprises two or more amino acids selected from the group consisting of glycine, serine, alanine, and threonine.
13. (Original) The method of claim 12, wherein step (b) is in excess of 30 µg per million cells in a 24 hour period.

14. (Withdrawn) ~~The method of claim 10, wherein the human IgG Fc variant comprises a hinge, CH2, and CH3 domains of human IgG2 with Pro331Ser mutation as SEQ ID NO: 18.~~
15. (Currently amended) The method of claim 10 14, wherein step (b) is in excess of 30 µg per million cells in a 24 hour period.
16. (Withdrawn)
17. (Withdrawn)
18. (Withdrawn)
19. (Withdrawn)
20. (Currently Amended) A method for making a recombinant fusion protein ~~comprising~~ consisting of HuEPO, a flexible peptide linker, and a human IgG Fc variant, which method comprises: (a) generating a CHO cell line transfected with DNA encoding the recombinant HuEPO-L-vFc fusion protein; (b) growing the cell line under conditions the recombinant protein is expressed in its growth medium in excess of 10 µg per million cells in a 24 hour period; and (c) purifying the expressed protein from step (b), wherein the recombinant fusion protein exhibits an enhanced *in vitro* biological activity of at least 2 fold relative to that of rHuEPO on a molar basis; wherein the flexible peptide linker containing about 20 or fewer amino acids is present between HuEPO and the human IgG Fc variant; and the peptide linker comprises two or more amino acids selected from the group consisting of glycine, serine, alanine, and threonine; wherein the human IgG Fc variant comprises a hinge, CH2, and CH3 domains selected from the group consisting of human IgG2 with Pro331Ser mutation as SEQ ID NO: 18, human IgG4 with Ser228Pro and Leu235Ala mutations as SEQ ID NO: 20, and human IgG1 with Leu234Val, Leu235Ala, and Pro331Ser mutations as SEQ ID NO: 22.